



## International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM) Codes for Xofigo

### ICD-10-CM codes associated with Xofigo:

ICD-10-CM Code <sup>1</sup>	Code Description
<b>C61</b>	Malignant neoplasm of prostate
<b>C79.51 or</b>	Secondary malignant neoplasm of bone
<b>C79.52</b>	Secondary malignant neoplasm of bone marrow

Information provided in this resource is for informational purposes only and does not guarantee that codes will be appropriate or that coverage and reimbursement will result. Customers should consult with their payers for all relevant coverage, coding, and reimbursement requirements. It is the sole responsibility of the provider to select proper codes and ensure the accuracy of all claims used in seeking reimbursement. Neither this resource nor Xofigo® Access Services is intended as legal advice or as a substitute for a provider's independent professional judgment.

### Contact Xofigo® Access Services Today for Reimbursement Support



**Phone:** 1-855-6XOFIGO (1-855-696-3446)



**Fax:** 1-855-963-4463



**Hours:** 9:00 AM – 7:00 PM ET,  
Monday through Friday



**Website:** [www.xofigo-us.com](http://www.xofigo-us.com)

**Provider Portal:**

<https://XofigoAccessOnline.com>

### INDICATION

Xofigo® is indicated for the treatment of patients with castration-resistant prostate cancer, symptomatic bone metastases and no known visceral metastatic disease.

### IMPORTANT SAFETY INFORMATION

**Contraindications:** Xofigo is contraindicated in women who are or may become pregnant. Xofigo can cause fetal harm when administered to a pregnant woman

Please see additional Important Safety Information on following page and [click here for full Prescribing Information.](#)

## IMPORTANT SAFETY INFORMATION (continued)

### Warnings and Precautions:

- **Bone Marrow Suppression:** In the phase 3 ALSYMPCA trial, 2% of patients in the Xofigo arm experienced bone marrow failure or ongoing pancytopenia, compared to no patients treated with placebo. There were two deaths due to bone marrow failure. For 7 of 13 patients treated with Xofigo bone marrow failure was ongoing at the time of death. Among the 13 patients who experienced bone marrow failure, 54% required blood transfusions. Four percent (4%) of patients in the Xofigo arm and 2% in the placebo arm permanently discontinued therapy due to bone marrow suppression. In the randomized trial, deaths related to vascular hemorrhage in association with myelosuppression were observed in 1% of Xofigo-treated patients compared to 0.3% of patients treated with placebo. The incidence of infection-related deaths (2%), serious infections (10%), and febrile neutropenia (<1%) was similar for patients treated with Xofigo and placebo. Myelosuppression—notably thrombocytopenia, neutropenia, pancytopenia, and leukopenia—has been reported in patients treated with Xofigo.

Monitor patients with evidence of compromised bone marrow reserve closely and provide supportive care measures when clinically indicated. Discontinue Xofigo in patients who experience life-threatening complications despite supportive care for bone marrow failure

- **Hematological Evaluation:** Monitor blood counts at baseline and prior to every dose of Xofigo. Prior to first administering Xofigo, the absolute neutrophil count (ANC) should be  $\geq 1.5 \times 10^9/L$ , the platelet count  $\geq 100 \times 10^9/L$ , and hemoglobin  $\geq 10$  g/dL. Prior to subsequent administrations, the ANC should be  $\geq 1 \times 10^9/L$  and the platelet count  $\geq 50 \times 10^9/L$ . Discontinue Xofigo if hematologic values do not recover within 6 to 8 weeks after the last administration despite receiving supportive care

- **Concomitant Use With Chemotherapy:** Safety and efficacy of concomitant chemotherapy with Xofigo have not been established. Outside of a clinical trial, concomitant use of Xofigo in patients on chemotherapy is not recommended due to the potential for additive myelosuppression. If chemotherapy, other systemic radioisotopes, or hemibody external radiotherapy are administered during the treatment period, Xofigo should be discontinued

- **Increased Fractures and Mortality in Combination With Abiraterone Plus Prednisone/Prednisolone:** Xofigo is not recommended for use in combination with abiraterone acetate plus prednisone/prednisolone outside of clinical trials. At the primary analysis of the phase 3 ERA-223 study that evaluated concurrent initiation of Xofigo in combination with abiraterone acetate plus prednisone/prednisolone in 806 asymptomatic or mildly symptomatic mCRPC patients, an increased incidence of fractures (28.6% vs 11.4%) and deaths (38.5% vs 35.5%) have been observed in patients who received Xofigo in combination with abiraterone acetate plus prednisone/prednisolone compared to patients who received placebo in combination with abiraterone acetate plus prednisone/prednisolone. Safety and efficacy with the combination of Xofigo and agents other than gonadotropin-releasing hormone analogues have not been established

**Administration and Radiation Protection:** Xofigo should be received, used, and administered only by authorized persons in designated clinical settings. The administration of Xofigo is associated with potential risks to other persons from radiation or contamination from spills of bodily fluids such as urine, feces, or vomit. Therefore, radiation protection precautions must be taken in accordance with national and local regulations

**Adverse Reactions:** The most common adverse reactions ( $\geq 10\%$ ) in the Xofigo arm vs the placebo arm, respectively, were nausea (36% vs 35%), diarrhea (25% vs 15%), vomiting (19% vs 14%), and peripheral edema (13% vs 10%). Grade 3 and 4 adverse events were reported in 57% of Xofigo-treated patients and 63% of placebo-treated patients. The most common hematologic laboratory abnormalities in the Xofigo arm ( $\geq 10\%$ ) vs the placebo arm, respectively, were anemia (93% vs 88%), lymphocytopenia (72% vs 53%), leukopenia (35% vs 10%), thrombocytopenia (31% vs 22%), and neutropenia (18% vs 5%)

Please click here for full [Prescribing Information](#).

1. 2014 International Classification of Diseases, 10th Revision, Clinical Modification Mappings. OptumInsights, Inc., 2013.



© 2018 Bayer. All rights reserved.  
BAYER, the Bayer Cross, Xofigo, and the Xofigo Access Services logo are registered trademarks of Bayer.  
PP-600-US-3660 09/18

