

## Available by prescription only.

Xofigo® (radium Ra 223 dichloride) injection is used to treat prostate cancer that no longer responds to hormonal or surgical treatment that lowers testosterone. It is for men whose prostate cancer has spread to the bone with symptoms but not to other parts of the body.



## Patient Treatment Guide

Your physician has referred you for Xofigo therapy. Your healthcare provider will give you a brochure that explains more about Xofigo. Over the course of your therapy, you will get up to 6 intravenous injections of Xofigo—1 injection every 4 weeks. Each injection of Xofigo will be given over 1 minute. This document will help you keep track of some of the important steps in your treatment. Before each injection of Xofigo, you will need to have a blood test so that your healthcare provider can be sure your blood counts are in the appropriate range. If your blood counts are not in the appropriate range, your scheduled treatment may be postponed or suspended.

### Injection 1

Your blood test will be drawn at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

Your first treatment will take place at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

### Injection 4

Your blood test will be drawn at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

Your fourth treatment will take place at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

### Injection 2

Your blood test will be drawn at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

Your second treatment will take place at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

### Injection 5

Your blood test will be drawn at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

Your fifth treatment will take place at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

### Injection 3

Your blood test will be drawn at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

Your third treatment will take place at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

### Injection 6

Your blood test will be drawn at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

Your sixth treatment will take place at (location) \_\_\_\_\_  
\_\_\_\_\_ on (date/time) \_\_\_\_\_

For questions call \_\_\_\_\_

These projected dates are specific to only your Xofigo treatment. While being treated with Xofigo, you will continue to see the healthcare provider who manages your cancer, just like you normally would. Please keep any additional appointments that your doctor requests.

Be sure to tell your healthcare provider about any signs of low blood cell counts. Signs of low blood cell counts include more bleeding or bruising than usual after injury, catching more colds than usual, fever, shortness of breath, and tiredness.

## IMPORTANT SAFETY INFORMATION

Xofigo is not indicated for women. Women who are pregnant or may become pregnant should not receive Xofigo and should practice safe sex, such as having their partner wear a condom and using effective birth control, as Xofigo can harm unborn babies. People who are handling fluids such as urine, feces, or vomit of a man taking Xofigo should wear gloves and wash their hands as precaution.



## IMPORTANT SAFETY INFORMATION (continued)

Before taking Xofigo, tell your healthcare provider if you:

- have bone marrow problems. Xofigo can cause your blood cells counts to go down, including red blood cells, white blood cells, and/or platelets. In a clinical trial, some patients had to permanently discontinue therapy because of bone marrow problems. In addition, there were some deaths and blood transfusions that occurred due to severe bone marrow problems. Your healthcare provider will do blood tests before and during treatment with Xofigo
- are receiving any chemotherapy or another extensive radiation therapy
- are taking Zytiga® (also known as abiraterone) with prednisone/prednisolone, as this may increase the risk of bone fractures (breaks) and death
- have any other medical conditions

While you are on Xofigo:

- make sure you keep your blood cell count monitoring appointments and tell your healthcare provider about any symptoms or signs of low blood cell counts. Report symptoms or signs of shortness of breath, tiredness, bleeding (such as bruising), or infection (such as fever)
- stay well hydrated and report any signs of dehydration (such as dry mouth and increased thirst), or urinary or kidney problems (such as burning when urinating)
- there are no restrictions regarding contact with other people after receiving Xofigo
- follow good hygiene practices in order to minimize radiation exposure from spills of bodily fluids to household members and caregivers for a period of one week after each injection
- use condoms and make sure female partners who may become pregnant use highly effective birth control methods during and for a minimum of six months after treatment with Xofigo

The most common side effects of Xofigo include:

- nausea
- diarrhea
- vomiting
- swelling of the arms or legs (peripheral edema)
- low blood cell counts

Tell your healthcare provider if you have any side effects that bother you or do not go away.

For important risk and use information about Xofigo, please click here for full [Prescribing Information](#).

You are encouraged to report negative side effects or quality complaints of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Xofigo is available by prescription only.  
For further information, please visit [www.xofigo-us.com](http://www.xofigo-us.com)

